



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>A61K 35/76, 48/00</b>	<b>A1</b>	(11) International Publication Number: <b>WO 97/12622</b> (43) International Publication Date: 10 April 1997 (10.04.97)
(21) International Application Number: PCT/US96/15406 (22) International Filing Date: 26 September 1996 (26.09.96) (30) Priority Data: 08/540,259 6 October 1995 (06.10.95) US (71) Applicant: THE SALK INSTITUTE FOR BIOLOGICAL STUDIES [US/US]; 10280 North Torrey Pines Road, La Jolla, CA 92037 (US). (72) Inventors: VERMA, Inder; 448 Marview Drive, Solana Beach, CA 92075 (US). TRONO, Didier; 457 Dell Court, San Diego, CA 92122 (US). NALDINI, Luigi; 13123 Shalimar Place, Del Mar, CA 92014 (US). GALLAY, Philippe; 930 Via Mil Cumbres #179, Solana Beach, CA 92075 (US). (74) Agent: HAILE, Lisa, A.; Fish & Richardson P.C., Suite 1400, 4225 Executive Square, La Jolla, CA 92037 (US).		(81) Designated States: AU, CA, IL, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>

(54) Title: VECTOR AND METHOD OF USE FOR NUCLEIC ACID DELIVERY TO NON-DIVIDING CELLS

## (57) Abstract

A recombinant retrovirus capable of infecting a non-dividing cell and a method of producing such a virus is provided. The recombinant retrovirus is preferably of lentivirus origin and is useful for the treatment of a variety of disorders including neurological disorders and disorders of other non-dividing cells.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic			SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

## VECTOR AND METHOD OF USE FOR NUCLEIC ACID DELIVERY TO NON-DIVIDING CELLS

This invention was made with Government support from grant no. RO1 AI37510 and CA 44360 awarded by the National Institutes of Health. The Government has certain  
5 rights in the invention.

### Field of the Invention

The present invention relates generally to the field of viral vectors and specifically to a novel recombinant retrovirus useful for the transfer and expression of nucleic acid sequences in non-dividing cells.

### Background of the Invention

10 The development of genetic vectors has heralded the fast-growing field of somatic gene transfer. Vectors based on simple retroviruses, such as the Moloney Leukemia Virus (MLV), are often selected because they efficiently integrate into the genome of the target cell. Integration is thought to be a prerequisite for long-term expression of  
15 the transduced gene. However, currently available retroviral vectors can only integrate in actively dividing cells which severely limits their use for *in vivo* gene transfer. Non-dividing cells are the predominant, long-lived cell type in the body, and account for most desirable targets of gene transfer, including liver, muscle, and brain. Even protocols attempting the transduction of hematopoietic stem cells require  
20 demanding *ex vivo* procedures for triggering cell division in these cells prior to infection.

In the early steps of infection, retroviruses deliver their nucleoprotein core into the cytoplasm of the target cell. Here, reverse transcription of the viral genome takes place while the core matures into a preintegration complex. The complex must reach  
25 the nucleus to achieve integration of the viral DNA into the host cell chromosomes. For simple retroviruses (oncoretroviruses), this step requires the dissolution of the nuclear membrane at mitotic prophase, most likely because the bulky size of the preintegration complex prevents its passive diffusion through the nuclear pores.

30 While retroviral vectors are useful for many kinds of *in vitro* gene transfer studies, problems including low titers limits their use for some *in vitro* and most *in vivo* studies. Further, another problem is that integration of retroviral vectors into the host genome was thought to be restricted to cells undergoing DNA replication. Thus, although

-2-

retroviral vectors capable of infecting a broad class of cell types are known, cell division is necessary for the proviral integration of these vectors. This effectively restricts the efficient use of retrovirus vectors to replicating cells. Thus, retroviruses have not been utilized to introduce genes into non-dividing or post-mitotic cells.

5 A possible way around this obstacle was suggested by recent studies of the pathogenesis of lentiviral diseases. Lentiviruses are complex retroviruses, which, in addition to the common retroviral genes gag, pol, and env, contain other genes with regulatory or structural function. The higher complexity enables the virus to modulate its life cycle, as in the course of latent infection. A typical lentivirus is the Human  
10 Immunodeficiency Virus (HIV), the etiologic agent of AIDS. *In vivo*, HIV can infect macrophages which are terminally differentiated cells that rarely divide. *In vitro*, HIV can infect primary cultures of monocyte-derived macrophages (MDM), and HeLa-Cd4 or T lymphoid cells arrested in the cell cycle by treatment with aphidicolin or  $\gamma$  irradiation. Infection of these cells is dependent on the active nuclear import of HIV  
15 preintegration complexes through the nuclear pores of the target cells. This occurs by the interaction of multiple, partly redundant, molecular determinants in the complex with the nuclear import machinery of the target cell. Identified determinants include a functional nuclear localization signal (NLS) in the gag MA protein, the karyophilic virion-associated protein vpr, and a C-terminal phosphotyrosine residue in a subset  
20 of the gag MA protein.

#### Summary of the Invention

The present invention provides a recombinant retrovirus that infects non-dividing host cells and transfers nucleic acid sequences which can then be expressed in the host cell. Thus, the present invention solves a long-felt need for a means of delivery of  
25 nucleic acid sequences to non-dividing cells.

In a first embodiment, the invention provides a recombinant retrovirus capable of infecting a non-dividing cell comprising a viral GAG, a viral POL, a viral ENV, a heterologous nucleic acid sequence operably linked to a regulatory nucleic acid sequence, and *cis*-acting nucleic acid sequences necessary for reverse transcription and integration. The recombinant retrovirus of the invention is preferably a lentivirus.  
30

In another embodiment, the invention provides a method of producing a recombinant retrovirus capable of infecting a non-dividing cell comprising transfecting a suitable

host cell with the following: a vector providing a nucleic acid encoding a viral gag and a viral pol, a vector providing a nucleic acid encoding a viral env, a vector providing a nucleic acid sequence encoding a packaging signal flanked by *cis*-acting nucleic acid sequences for reverse transcription and integration, and providing a cloning site  
5 for introduction of a heterologous nucleic acid sequence, operably linked to a regulatory nucleic acid sequence, and recovering the recombinant virus.

In yet another embodiment, the invention provides a method for introduction and expression of a heterologous nucleic acid sequence in a non-dividing cell comprising  
10 infecting the non-dividing cell with the recombinant virus of the invention and expressing the heterologous nucleic acid sequence in the non-dividing cell.

#### **Brief Description of the Drawings**

Figure 1 is a schematic illustration of the method for production of an HIV-based recombinant retrovirus.

Figure 2 is a table of the titer of infectious virus particles using a VSV.G env, LacZ  
15 reporter gene and a packaging construct having an HIV (pCMVDR8) or MLV gag/pol before and after concentration (= "yield").

Figure 3 is a table of titers of recombinant virus after co-transfection of 293T cells (fibroblast packaging cell line) with HIV-based and MLV-based packaging constructs.

Figure 4 is a graph of the efficiency of transduction of HIV and MLV-based CMV- $\beta$ -galactosidase vectors into growing HeLa cells, cells arrested at the G1/S interphase  
20 by aphidicolin, and cells arrested at G2 by x-rays.

Figure 5 is a graph of the efficiency of transduction of HIV and MLV CMV-luciferase vectors into rat 208F fibroblasts in G0 after 4, 7, 11, and 15 days.

Figure 6 is a graph of the efficiency of transduction of HIV and MLV CMV-luciferase  
25 vectors into human primary macrophages.

Figure 7 is a graph of survival of HIV- and MLV-based vector in cells arrested for 3 weeks, inoculated with the vectors, and induced to divide at indicated times after infection.

### Description of the Preferred Embodiments

The present invention provides a recombinant retrovirus capable of infecting non-dividing cells. The virus is useful for the *in vivo* and *ex vivo* transfer and expression of genes nucleic acid sequences (e.g., in non-dividing cells).

5 Retroviruses are RNA viruses wherein the viral genome is RNA. When a host cell is infected with a retrovirus, the genomic RNA is reverse transcribed into a DNA intermediate which is integrated very efficiently into the chromosomal DNA of infected cells. This integrated DNA intermediate is referred to as a provirus. Transcription of  
10 the provirus and assembly into infectious virus occurs in the presence of an appropriate helper virus or in a cell line containing appropriate sequences enabling encapsidation without coincident production of a contaminating helper virus. As described below, a helper virus is not required for the production of the recombinant retrovirus of the present invention, since the sequences for encapsidation are provided by co-transfection with appropriate vectors.

15 The retroviral genome and the proviral DNA have three genes: the *gag*, the *pol*, and the *env*, which are flanked by two long terminal repeat (LTR) sequences. The *gag* gene encodes the internal structural (matrix, capsid, and nucleocapsid) proteins; the *pol* gene encodes the RNA-directed DNA polymerase (reverse transcriptase) and the *env* gene encodes viral envelope glycoproteins. The 5' and 3' LTRs serve to promote  
20 transcription and polyadenylation of the virion RNAs. The LTR contains all other *cis*-acting sequences necessary for viral replication. Lentiviruses have additional genes including *vif*, *vpr*, *tat*, *rev*, *vpu*, *nef*, and *vpx* (in HIV-1, HIV-2 and/or SIV).

25 Adjacent to the 5' LTR are sequences necessary for reverse transcription of the genome (the tRNA primer binding site) and for efficient encapsidation of viral RNA into particles (the Psi site). If the sequences necessary for encapsidation (or packaging of retroviral RNA into infectious virions) are missing from the viral genome, the result is a *cis* defect which prevents encapsidation of genomic RNA. However, the resulting mutant is still capable of directing the synthesis of all virion  
30 proteins.

In a first embodiment, the invention provides a recombinant retrovirus capable of infecting a non-dividing cell. The recombinant retrovirus comprises a viral GAG, a viral POL, a viral ENV, a heterologous nucleic acid sequence operably linked to a

regulatory nucleic acid sequence, and *cis*-acting nucleic acid sequences necessary for packaging, reverse transcription and integration, as described above. It should be understood that the recombinant retrovirus of the invention is capable of infecting dividing cells as well as non-dividing cells.

5 The recombinant retrovirus of the invention is therefore genetically modified in such a way that some of the structural, infectious genes of the native virus have been removed and replaced instead with a nucleic acid sequence to be delivered to a target non-dividing cell. After infection of a cell by the virus, the virus injects its  
10 nucleic acid into the cell and the retrovirus genetic material can integrate into the host cell genome. The transferred retrovirus genetic material is then transcribed and translated into proteins within the host cell.

The invention provides a method of producing a recombinant retrovirus capable of infecting a non-dividing cell comprising transfecting a suitable host cell with the following: a vector providing a nucleic acid encoding a viral gag and a viral pol; a  
15 vector providing a nucleic acid encoding a viral env; a vector providing a nucleic acid sequence encoding a packaging signal flanked by *cis*-acting nucleic acid sequences for reverse transcription and integration, and providing a cloning site for introduction of a heterologous gene, operably linked to a regulatory nucleic acid sequence, and recovering the recombinant virus. An illustration of the individual vectors used in the  
20 method of the invention is shown in FIGURE 1.

The method of the invention includes the combination of a minimum of three vectors in order to produce a recombinant virion or recombinant retrovirus. A first vector provides a nucleic acid encoding a viral gag and a viral pol. (see the illustrative  
25 Packaging Construct, FIGURE 1) These sequences encode a group specific antigen and reverse transcriptase, (and integrase and protease-enzymes necessary for maturation and reverse transcription), respectively, as discussed above. Most preferably, the viral gag and pol are derived from a lentivirus, and most preferably from HIV.

30 A second vector provides a nucleic acid encoding a viral envelope (env). The env gene can be derived from any virus, including retroviruses. The env may be amphotropic envelope protein which allows transduction of cells of human and other species, or may be ecotropic envelope protein, which is able to transduce only mouse and rat cells. Further, it may be desirable to target the recombinant virus by linkage

of the envelope protein with an antibody or a particular ligand for targeting to a receptor of a particular cell-type. By inserting a sequence (including regulatory region) of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific. Retroviral vectors can be made target specific by inserting, for example, a glycolipid, or a protein. Targeting is often accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific methods to achieve delivery of a retroviral vector to a specific target.

Examples of retroviral-derived env genes include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), gibbon ape leukemia virus (GaLV), human immunodeficiency virus (HIV) and Rous Sarcoma Virus (RSV). Other env genes such as Vesicular stomatitis virus (VSV) (Protein G) can also be used.

The vector providing the viral env nucleic acid sequence is operably associated with regulatory sequence, e.g., a promoter or enhancer. (see Pseudotyping ENV plasmid, FIGURE 1) Preferably, the regulatory sequence is a viral promoter. The regulatory sequence can be any eukaryotic promoter or enhancer, including for example, the Moloney murine leukemia virus promoter-enhancer element, the human cytomegalovirus enhancer (as used in the illustrative example), or the vaccinia P7.5 promoter. In some cases, such as the Moloney murine leukemia virus promoter-enhancer element, these promoter-enhancer elements are located within or adjacent to the LTR sequences.

A third vector provides the cis-acting viral sequences necessary for the viral life cycle. Such sequences include the  $\psi$  packaging sequence, reverse transcription signals, integration signals, viral promoter, enhancer, and polyadenylation sequences. The third vector also contains a cloning site for a heterologous nucleic acid sequence to be transferred to a non-dividing cell. A schematic illustration of a suitable vector is shown in FIGURE 1).

Since recombinant retroviruses produced by standard methods in the art are defective, they require assistance in order to produce infectious vector particles. Typically, this assistance is provided, for example, by using a helper cell line that provides the missing viral functions. These plasmids are missing a nucleotide



-7-

sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include but are not limited to  $\Psi$ 2, PA317 and PA12, for example. Suitable cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced.

The method of producing the recombinant retrovirus of the invention is different than the standard helper virus/packaging cell line method described above. The three or more individual vectors used to co-transfect a suitable packaging cell line collectively contain all of the required genes for production of a recombinant virus for infection and transfer of nucleic acid to a non-dividing cell. Consequently, there is no need for a helper virus.

The heterologous nucleic acid sequence is operably linked to a regulatory nucleic acid sequence. As used herein, the term "heterologous" nucleic acid sequence refers to a sequence that originates from a foreign species, or, if from the same species, it may be substantially modified from its original form. Alternatively, an unchanged nucleic acid sequence that is not normally expressed in a cell is a heterologous nucleic acid sequence. The term "operably linked" refers to functional linkage between the regulatory sequence and the heterologous nucleic acid sequence. Preferably, the heterologous sequence is linked to a promoter, resulting in a chimeric gene. The heterologous nucleic acid sequence is preferably under control of either the viral LTR promoter-enhancer signals or of an internal promoter, and retained signals within the retroviral LTR can still bring about efficient integration of the vector into the host cell genome.

The promoter sequence may be homologous or heterologous to the desired gene sequence. A wide range of promoters may be utilized, including viral or mammalian promoter. Cell or tissue specific promoters can be utilized to target expression of gene sequences in specific cell populations. Suitable mammalian and viral promoters for the present invention are available in the art.

Preferably during the cloning stage, the nucleic acid construct referred to as the transfer vector, having the packaging signal and the heterologous cloning site, also contains a selectable marker gene. Marker genes are utilized to assay for the

presence of the vector, and thus, to confirm infection and integration. The presence of this marker gene ensures the growth of only those host cells which express the inserts. Typical selection genes encode proteins that confer resistance to antibiotics and other toxic substances, e.g., histidinol, puromycin, hygromycin, neomycin, methotrexate, etc. The illustrative examples herein utilize the  $\beta$ -galactosidase (LacZ) or luciferase reporter or marker system.

The recombinant virus of the invention is capable of transferring a nucleic acid sequence into a non-dividing cell. The term nucleic acid sequence refers to any nucleic acid molecule, preferably DNA. The nucleic acid molecule may be derived from a variety of sources, including DNA, cDNA, synthetic DNA, RNA, or combinations thereof. Such nucleic acid sequences may comprise genomic DNA which may or may not include naturally occurring introns. Moreover, such genomic DNA may be obtained in association with promoter regions, introns, or poly A sequences. Genomic DNA may be extracted and purified from suitable cells by means well known in the art. Alternatively, messenger RNA (mRNA) can be isolated from cells and used to produce cDNA by reverse transcription or other means.

The phrase "non-dividing" cell refers to a cell that does not go through mitosis. Non-dividing cells may be blocked at any point in the cell cycle, (e.g., G<sub>0</sub>/G<sub>1</sub>, G<sub>1</sub>/S, G<sub>2</sub>/M), as long as the cell is not actively dividing. For *ex vivo* infection, a dividing cell can be treated to block cell division by standard techniques used by those of skill in the art, including, irradiation, aphidocolin treatment, serum starvation, and contact inhibition. However, it should be understood that *ex vivo* infection is often performed without blocking the cells since many cells are already arrested (e.g., stem cells). The recombinant retrovirus vector of the invention is capable of infecting any non-dividing cell, regardless of the mechanism used to block cell division or the point in the cell cycle at which the cell is blocked. Examples of pre-existing non-dividing cells in the body include neuronal, muscle, liver, skin, heart, lung, and bone marrow cells, and their derivatives.

Preferably, the recombinant retrovirus produced by the method of the invention is lentivirus-derived, and more preferably the recombinant lentivirus is a derivative of human immunodeficiency virus (HIV). For public policy reasons, typically when the retrovirus is HIV, the *env* will be derived from a virus other than HIV.

The method of the invention provides at least three vectors which provide all of the functions required for packaging of recombinant virions as discussed above. The method also envisions transfection of vectors including viral genes such as vpr, vif, nef, vpx, tat, rev, and vpu. Some or all of these genes can be included on the packaging construct vector, for example, or, alternatively, they may reside on  
5 individual vectors. There is no limitation to the number of vectors which are utilized, as long as they are co-transfected to the packaging cell line in order to produce a single recombinant retrovirus. For example, one could put the env nucleic acid sequence on the same construct as the gag and pol.

10 The vectors are introduced via transfection or infection into the packaging cell line. The packaging cell line produces viral particles that contain the vector genome. Methods for transfection or infection are well known by those of skill in the art. After co-transfection of the at least three vectors to the packaging cell line, the recombinant virus is recovered from the culture media and titered by standard methods used by  
15 those of skill in the art.

In another embodiment, the invention provides a recombinant retrovirus produced by the method of the invention as described above.

The invention also provides a recombinant retrovirus capable of infecting a non-dividing cell comprising a viral GAG; a viral POL; a viral ENV; a heterologous nucleic acid sequence operably linked to a regulatory nucleic acid sequence; and cis-acting  
20 nucleic acid sequences necessary for packaging, reverse transcription and integration. The recombinant retrovirus is preferably a lentivirus, and most preferably HIV. As described above for the method of producing a recombinant retrovirus above, the recombinant retrovirus of the invention may further include at least one of VPR,  
25 VIF, NEF, VPX, TAT, REV, and VPU protein. While not wanting to be bound by a particular theory, it is believed that one or more of these genes/protein products are important for increasing the viral titer of the recombinant retrovirus produced (e.g., NEF) or may be necessary for infection and packaging of virions, depending on the packaging cell line chosen (e.g., VIF).

30 The invention also provides a method of nucleic acid transfer to a non-dividing cell to provide expression of a particular nucleic acid sequence. Therefore, in another embodiment, the invention provides a method for introduction and expression of a heterologous nucleic acid sequence in a non-dividing cell comprising infecting the

-10-

non-dividing cell with the recombinant virus of the invention and expressing the heterologous nucleic acid sequence in the non-dividing cell.

It may be desirable to modulate the expression of a gene regulating molecule in a cell by the introduction of a molecule by the method of the invention. The term "modulate" envisions the suppression of expression of a gene when it is over-expressed, or augmentation of expression when it is under-expressed. Where a cell proliferative disorder is associated with the expression of a gene, nucleic acid sequences that interfere with the gene's expression at the translational level can be used. This approach utilizes, for example, antisense nucleic acid, ribozymes, or triplex agents to block transcription or translation of a specific mRNA, either by masking that mRNA with an antisense nucleic acid or triplex agent, or by cleaving it with a ribozyme.

Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule (Weintraub, *Scientific American*, 262:40, 1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the mRNA, since the cell will not translate a mRNA that is double-stranded. Antisense oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to cause problems than larger molecules when introduced into the target cell. The use of antisense methods to inhibit the *in vitro* translation of genes is well known in the art (Marcus-Sakura, *Anal. Biochem.*, 172:289, 1988).

The antisense nucleic acid can be used to block expression of a mutant protein or a dominantly active gene product, such as amyloid precursor protein that accumulates in Alzheimer's disease. Such methods are also useful for the treatment of Huntington's disease, hereditary Parkinsonism, and other diseases. Antisense nucleic acids are also useful for the inhibition of expression of proteins associated with toxicity.

Use of an oligonucleotide to stall transcription is known as the triplex strategy since the oligomer winds around double-helical DNA, forming a three-strand helix. Therefore, these triplex compounds can be designed to recognize a unique site on a chosen gene (Maher, *et al.*, *Antisense Res. and Dev.*, 1(3):227, 1991; Helene, C., *Anticancer Drug Design*, 6(6):569, 1991).

Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA in a manner analogous to DNA restriction endonucleases. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific nucleotide sequences in an RNA molecule and cleave it (Cech, *J.Amer.Med. Assn.*, 260:3030, 1988). A major advantage of this approach is that, because they are sequence-specific, only mRNAs with particular sequences are inactivated.

It may be desirable to transfer a nucleic acid encoding a biological response modifier. Included in this category are immunopotentiating agents including nucleic acids encoding a number of the cytokines classified as "interleukins". These include, for example, interleukins 1 through 12. Also included in this category, although not necessarily working according to the same mechanisms, are interferons, and in particular gamma interferon ( $\gamma$ -IFN), tumor necrosis factor (TNF) and granulocyte-macrophage-colony stimulating factor (GM-CSF). It may be desirable to deliver such nucleic acids to bone marrow cells or macrophages to treat enzymatic deficiencies or immune defects. Nucleic acids encoding growth factors, toxic peptides, ligands, receptors, or other physiologically important proteins can also be introduced into specific non-dividing cells.

The recombinant retrovirus of the invention can be used to treat an HIV infected cell (e.g., T-cell or macrophage) with an anti-HIV molecule. In addition, respiratory epithelium, for example, can be infected with a recombinant retrovirus of the invention having a gene for cystic fibrosis transmembrane conductance regulator (CFTR) for treatment of cystic fibrosis.

The method of the invention may also be useful for neuronal or glial cell transplantation, or "grafting", which involves transplantation of cells infected with the recombinant retrovirus of the invention *ex vivo*, or infection *in vivo* into the central nervous system or into the ventricular cavities or subdurally onto the surface of a host brain. Such methods for grafting will be known to those skilled in the art and are described in *Neural Grafting in the Mammalian CNS*, Bjorklund and Stenevi, eds., (1985), incorporated by reference herein. Procedures include intraparenchymal transplantation, (i.e., within the host brain) achieved by injection or deposition of tissue within the host brain so as to be apposed to the brain parenchyma at the time of transplantation.

-12-

Administration of the cells or virus into selected regions of the recipient subject's brain may be made by drilling a hole and piercing the dura to permit the needle of a microsyringe to be inserted. The cells or recombinant retrovirus can alternatively be injected intrathecally into the spinal cord region. A cell preparation infected *ex vivo*,  
5 or the recombinant retrovirus of the invention, permits grafting of neuronal cells to any predetermined site in the brain or spinal cord, and allows multiple grafting simultaneously in several different sites using the same cell suspension or viral suspension and permits mixtures of cells from different anatomical regions.

Cells infected with a recombinant retrovirus of the invention, *in vivo*, or *ex vivo*, used  
10 for treatment of a neuronal disorder for example, may optionally contain an exogenous gene, for example, a gene which encodes a receptor or a gene which encodes a ligand. Such receptors include receptors which respond to dopamine, GABA, adrenaline, noradrenaline, serotonin, glutamate, acetylcholine and other neuropeptides, as described above. Examples of ligands which may provide a  
15 therapeutic effect in a neuronal disorder include dopamine, adrenaline, noradrenaline, acetylcholine, gamma-aminobutyric acid and serotonin. The diffusion and uptake of a required ligand after secretion by an infected donor cell would be beneficial in a disorder where the subject's neural cell is defective in the production of such a gene product. A cell genetically modified to secrete a neurotrophic factor, such as nerve  
20 growth factor, (NGF), might be used to prevent degeneration of cholinergic neurons that might otherwise die without treatment. Alternatively, cells be grafted into a subject with a disorder of the basal ganglia, such as Parkinson's disease, can be modified to contain an exogenous gene encoding L-DOPA, the precursor to dopamine. Parkinson's disease is characterized by a loss of dopamine neurons in the  
25 substantia-nigra of the midbrain, which have the basal ganglia as their major target organ.

Other neuronal disorders that can be treated similarly by the method of the invention include Alzheimer's disease, Huntington's disease, neuronal damage due to stroke, and damage in the spinal cord. Alzheimer's disease is characterized by degeneration  
30 of the cholinergic neurons of the basal forebrain. The neurotransmitter for these neurons is acetylcholine, which is necessary for their survival. Engraftment of cholinergic cells infected with a recombinant retrovirus of the invention containing an exogenous gene for a factor which would promote survival of these neurons can be accomplished by the method of the invention, as described. Following a stroke, there  
35 is selective loss of cells in the CA1 of the hippocampus as well as cortical cell loss

which may underlie cognitive function and memory loss in these patients. Once identified, molecules responsible for CA1 cell death can be inhibited by the methods of this invention. For example, antisense sequences, or a gene encoding an antagonist can be transferred to a neuronal cell and implanted into the hippocampal region of the brain.

The method of transferring nucleic acid also contemplates the grafting of neuroblasts in combination with other therapeutic procedures useful in the treatment of disorders of the CNS. For example, the retroviral infected cells can be co-administered with agents such as growth factors, gangliosides, antibiotics, neurotransmitters, neurohormones, toxins, neurite promoting molecules and antimetabolites and precursors of these molecules such as the precursor of dopamine, L-DOPA.

Further, there are a number of inherited neurologic diseases in which defective genes may be replaced including: lysosomal storage diseases such as those involving  $\beta$ -hexosaminidase or glucocerebrosidase; deficiencies in hypoxanthine phosphoribosyl transferase activity (the "Lesch-Nyhan" syndrome); amyloid polyneuropathies (-prealbumin); Duchenne's muscular dystrophy, and retinoblastoma, for example.

For diseases due to deficiency of a protein product, gene transfer could introduce a normal gene into the affected tissues for replacement therapy, as well as to create animal models for the disease using antisense mutations. For example, it may be desirable to insert a Factor IX encoding nucleic acid into a retrovirus for infection of a muscle of liver cell.

The following Examples are intended to illustrate, but not to limit the invention. While such Examples are typical of those that might be used, other procedures known to those skilled in the art may alternatively be utilized.

### EXAMPLES

The following examples illustrate the three-plasmid lentiviral vector system of the invention. A non infectious plasmid provides *in trans* the structural and regulatory proteins of HIV but the envelope (packaging construct, pCMV $\Delta$ R8). The transducing vector contains all known *cis*-acting sequences of HIV-and cloning sites for the introduction of the gene to be transduced (vector plasmid, pHR'). In the experiments described hereafter, two marker genes (*E. coli*  $\beta$ -galactosidase -LacZ-, and firefly

-14-

luciferase -Lucif-), driven by the human Cytomegalovirus immediate early enhancer/promoter -CMV or C- have been cloned in the vector plasmid, thus generating pHR'CLacZ and pHR'-Clucif (FIGURE 1).

5 A third plasmid encodes a heterologous envelope to pseudotype the viral particles. Pseudotyping both broadens the host range of the vector, and increases its biosafety. Three types of env-encoding plasmid have been used in the present examples. Plasmid pSV-A-MLV-env (Page, *et al.*, 1990, *J. Virol.*, 64:5270, 1990) encodes the amphotropic envelope of the 4070 Moloney Leukemia Virus (MLV/Ampho) under the transcriptional control of the MLV LTR. Plasmid pCMV-Eco env encodes the ecotropic  
10 envelope of the Moloney Leukemia Virus (MLV/Eco) under the transcriptional control of the CMV promoter (N. Somia, Salk Institute, La Jolla, CA). Plasmid pMD.G encodes the envelope protein G of the Vesicular Stomatitis Virus (VSV.G) under the transcriptional control of the CMV promoter (D. Ory and R. Mulligan, Whitehead Institute, Cambridge, MA).

15 Replication-defective viral particles were produced by the transient co-transfection of the three plasmids in 293T human kidney cells. The conditioned medium was harvested, filtered, and assayed for the transduction of the reporter genes into target cells by standard methods. Human HeLa cells and rat 208F fibroblasts were tested in different growth conditions, and the dependence of the transduction on the  
20 progression along the cell cycle was evaluated. Growth arrest by density-dependent inhibition of growth (GO) was compared with pharmacological or other means of the cell cycle arrest (later phases of the cycle). The transduction of non-dividing, terminally differentiated cells such as macrophages and neurons was also tested, both *in vitro* and *in vivo*.

### 25 Example 1

#### CONSTRUCTION OF pCMVDR8

The packaging plasmid pCMVDR8 was constructed in a series of steps starting from the plasmid pR8, an infectious molecular clone of proviral HIV DNA. pR8 was made  
30 by replacing a BamHI-BssHII fragment in the plasmid pR7 (Kim *et al.*, *J. Virol.*, 63:3708, 1989; von Schwedler *et al.*, *J. Virol.*, 67:4945, 1993), which contains the HIV-1 HXB2d proviral DNA (Shaw *et al.*, *Science*, 226:1165, 1984) with a full-length nef reading frame, by the homologous fragment from the NL4.3 HIV proviral DNA. NL4.3 is a molecularly constructed provirus that contains full-length, functional vpr



-15-

and vpu reading frames (Adachi *et al.*, *J. Virol.*, 59:284, 1986). The backbone of plasmid pR7/R8 contains the Amp resistance gene and the pUC origin of replication from plasmid sp65, and the SV40 origin of replication and the gpt gene from plasmid pSVgpt. An env-defective version of pR8, pΔER8, was made by replacing the 2.7 kbp  
5 Sall-BamHI fragment with the homologous fragment from pΔER7, which contains a MluI linker inserted at the StuI site 6831 that frameshifts the env reading frame (Trono, *et al.*, *Cell.*, 59:113, 1989). A 39 bp internal deletion in the Ψ sequence between the 5' splice donor site and the starting codon of the gag gene, was introduced in pΔER8 by replacing a 797 bp BssHII-SpeI fragment with an homologous  
10 758 fragment containing a previously described deletion in the packaging signal (Aldovini & Young, *J. Virol.*, 64:1920, 1990). The resulting plasmid was called pΔΨΔER8. Plasmid pAΔΨΔER8 was constructed by replacing the 3' HIV LTR with the polyadenylation site of insulin genomic DNA (Trono, *et al.*, *Cell.*, 59:113, 1989) maintaining intact the nef reading frame. It was constructed by PCR, introducing a  
15 unique NotI site at the junction between the nef reading frame and the polyA signal.

In a first series of amplification, a 961 bp fragment was amplified with the oligonucleotides gp41.4 (s), overlapping the BamHI site in gp41 of HIV1HXB2- and nef27 (a) from the pΔER8 template, and a 402 bp fragment was amplified with the o-  
ligonucleotides LN-A (s) and LN-B (a) containing an XbaI site - from a DNA template  
20 containing the polyadenylation site of the insulin gene. Oligonucleotides nef27(a) and LN-A(s) contain a complementary sequence carrying a NotI restriction site. The two amplicates were purified, mixed and used as template for a second amplification with the oligonucleotides gp41.4(s) and LN-B(a) (see Table 1 for oligo sequences). The  
final 1.3 kbp product was purified, digested with BamHI and XbaI and cloned in the  
25 13.5 kbp BamHI-XbaI fragment of pΔΨΔER8. Plasmid pCMVΔ8 was constructed substituting the 5' LTR and leader sequences in pAΔΨΔER8 with the CMV promoter. A 0.8 kbp fragment containing the CMV promoter was lifted from the plasmid pCMVpA (W.W. Chang, University of Texas, M.D. Anderson Cancer Center, Houston, TX) by double digestion with SmaI and HpaI. A MluI linker was added to the HpaI  
30 ends and the fragment was cloned into the 14.5 kbp BssHII-HpaI fragment of pAΔΨΔER8. One of skill in the art will be able to identify similar starting plasmids and sequences without undue experimentation based on the teachings herein.

-16-

Table 1

<u>Oligonucleotide</u>	<u>Nucleotide sequence</u>
gp41.4 (s)	5' GTGAACGGATCCTTGGAACATC 3' (SEQ ID NO:1)
5    nef 27 (a)	5' CGGGGCGGCCGCTCAGCAGTTCT TGAAGTACTC 3' (SEQ ID NO:2)
LN-A (s)	5'CCGGAGTACTTCAAGAACTGCTGA GCGGCCGCCCCGGTGACCTTCAGA CCTTGGC 3' (SEQ ID NO:3)
10	
LN-B (a)	5' GCGGTCTAGACTAAGAAACCATTATTA TCATGAC 3' (SEQ ID NO:4)
gag31 (s)	5' AAGACCACCGCACAGCAAGCGGC CGCTGACACAGGACACAGCAAT 3' (SEQ ID NO:5)
15	

Example 2CONSTRUCTION OF pHR'

The vector plasmid pHR' was constructed by cloning a fragment of the env gene encompassing the RRE (responsive element) and a splice acceptor site between the two LTR's of the HIV-1 HXB2 proviral DNA. The leader and  $\Psi$  sequences were retained in the construct, together with the 5' 0.3 kbp of the gag gene, previously shown to constitute an extended packaging signal. A 854 bp env fragment (BglII 7620-BamHI 8474) was obtained from pR7. A NotI linker was added to the filled-in BglII end. The fragment was cloned into the 8.9 kbp NotI-BamHI fragment of plasmid pMAKK<sup>113</sup>TTR7, a pR7 derivative containing mutations in the gag codons 113, 114, and 118 (the last ones introduce a unique NotI cloning site; sequence as in oligonucleotide Gag 31 (s)). Plasmid pHR was obtained. Plasmid pHR' was generated introducing a frameshift mutation 42 bp downstream the gag gene AUG, which closes the reading frame of the gene fragment retained in the construct. The mutation was introduced by opening and filling-in a ClaI site. Two unique BamHI and XhoI sites are available in pHR' upstream of the 3' LTR for cloning inserts.

-17-

**Example 3****CONSTRUCTION OF pHR'-CLacZ and pHR'-Clucif**

5 A Sall-XhoI 3.6 kbp fragment containing the *E. coli*  $\beta$ -galactosidase gene under the transcriptional control of the CMV promoter was obtained from the plasmid pSLX-CMV-LacZ (Scharfmann *et al.*, *Proc. Natl. Acad. Sci. USA*, **88**:4626, 1991). After filling-in the Sall ends, the fragment was cloned into the 8.9 kbp XhoI-BamHI fragment of pHR', whose BamHI ends had previously been filled-in. pHR'-CLacZ was obtained. pHR'-CLucif was obtained replacing a 3.1 kbp BamHI-XhoI fragment in pHR'-CLacZ, containing the LacZ reading frame, with a 1.7 kbp BamHI-XhoI fragment from pGEM-luc (Promega) containing the firefly luciferase reading frame.

10

**Example 4****PRODUCTION OF VECTOR PARTICLES**

15 Replication-defective viral particles were produced by the transient cotransfection of the above-described plasmids in 293T human kidney cells. All plasmids were transformed and grown in *E. coli* HB101 bacteria following standard molecular biology procedures. For transfection of eukaryotic cells, plasmid DNA was purified twice by equilibrium centrifugation in CsCl-ethidium bromide gradients. A total of 40  $\mu$ g DNA was used for the transfection of a culture in a 10 cm dish, in the following proportions: 10  $\mu$ g pCMV $\Delta$ R8, 20  $\mu$ g pHR', and 10  $\mu$ g env plasmids, either MLV/Ampho, MLV/Eco or VSV.G. When a plasmid was omitted, or added in different proportion, pGEM-LacZ was added to maintain constant the total amount of DNA added to the cells. 293T cells were grown in DMEM supplemented with 10% fetal calf serum and antibiotics in a 10% CO<sub>2</sub> incubator. Cells were plated at a density of  $1.3 \times 10^6$ /10 cm dish the day before transfection. Culture medium was changed 4 to 6 hrs before transfection.

20 Calcium phosphate-DNA complexes were prepared according to the method of Chen and Okayama (*Mol. Cell. Biol.*, **7**:2745, 1987), and incubated overnight with the cells in an atmosphere of 5% CO<sub>2</sub>. The following morning, the medium was replaced, and the cultures returned to 10% CO<sub>2</sub>. Conditioned medium was harvested 48 to 60 hrs after transfection, cleared of cellular debris by low speed centrifugation (300 x g 10 min), and filtered through 0.45  $\mu$ m low protein binding filters. The conditioned medium was either used immediately for infection or kept frozen at -80°C. The content of viral particles was monitored by p24 gag antigen ELISA (DuPont).

25

30

-18-

MLV-based vectors were also produced from transient cotransfection in 293T cells of the following plasmids. Plasmid pSLX-CMVlacZ (Scharfmann *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:4626, 1991) is a MLV-derived vector carrying a CMV-driven *E. coli*  $\beta$ -galactosidase gene. The pCL plasmid series carry a hybrid CMV-LTR promoter which allows for CMV-driven transcription in the packaging cell and reconstitution of a functional LTR in the target cell (R. Naviaux, Salk Institute, La Jolla, CA). Plasmid pCLNC-Lucif is a MLV-derived vector carrying a CMV-driven Firefly luciferase gene. It was made by cloning a 1.7 kbp HindIII-StuI fragment from pGEM-luc (Promega), containing the luciferase reading frame, into the 8 kbp ClaI-HindIII fragment of pCLNCX, after filling-in the ClaI ends. Plasmid pCL-ECO is a ecotropic packaging plasmid. Plasmid pCMV-GAGPOL (N. Somia, Salk Institute, La Jolla, CA) is a semi-packaging plasmid carrying a CMV-driven gag-pol gene from MLV. It was used in conjunction with pMD.G to package VSV.G-pseudotyped MLV-based vectors.

Figure 2 shows a comparison between the infectious titers of recombinant lentivirus (HIV) construct and an MLV construct assayed on 208F fibroblasts. The results indicate that the HIV vector of the present invention is at least as infectious as the standard MLV vector. Purified stock indicated that titers after concentrating the supernatant by ultracentrifugation. Yield reflects percent recovery of virus after concentration.

Figure 3 shows the results of co-transfection of a packaging construct, env-encoding construct and transfer vector (as in Figure 1). Various HIV and MLV-based packaging constructs were compared using amphotropic and ecotropic env genes, as well as VSV.G env. The figure shows a comparison of infection of rat 208F cells with the resulting recombinant retroviruses. The titers for the HIV constructs having an MLV (amphotropic) or VSV.G env were similar to the standard MLV-based vector (approximately  $10^5$ ). The infectious titer for the HIV-based construct having an MLV ecotropic env was about 10-fold less than the amphotropic env (approx.  $10^4$ ).

**Example 5****CONCENTRATION OF VIRAL PARTICLES**

5 Pooled conditioned medium harvested as described above was layered on top of a cushion of 20% sucrose solution in PBS and centrifuged in a Beckman SW28 rotor at 50,000 x g for 90 min. The pellet was resuspended by incubation and gentle pipetting in 1-4 ml PBS for 30-60 min, then centrifuged again at 50,000 x g x 90 min in a Beckmann SW55 rotor. The pellet was resuspended in a minimal volume (20-50  $\mu$ l) of PBS and either used directly for infection or stored in frozen aliquots at -80°C.

**Example 6****INFECTION OF CULTURED CELLS**

10 For the infection of cultured cells, serial dilutions of conditioned medium from 293T transient transfectants or of concentrated viral stocks were added to the culture medium supplemented with 8  $\mu$ g/ml polybrene. The cells were incubated from 3 hrs to overnight, then the medium was replaced, and the cells further incubated for 36 hrs prior to assaying expression of the transduced gene. Expression of  $\beta$ -Galactosidase ( $\beta$ -Gal) was scored by fixing the cultures 5 min in cold 2% formaldehyde/0.2% glutaraldehyde in PBS, washing twice with PBS, and staining with 1 mg/ml X-gal in PBS containing 5 mM each of K ferri- and ferro-cyanide and 2 mM  $MgCl_2$ , for 4 hrs to overnight at 37°C. Titers were calculated counting the number of individual foci of blue cells per well, and multiplying it by the volume of vector stock added and the dilution factor. Expression of luciferase was assayed by washing the cultures twice with TBS and extracting the cells with 200  $\mu$ l/well 0.5% NP40 in TBS containing 5 mM  $MgCl_2$ . The extract was cleared by centrifugation at 15,000 x g for 10 min. 50  $\mu$ l of extract were diluted with 150  $\mu$ l of 75 mM Tris-HCl pH=7.8, 15 mM Mg acetate and 4 mM ATP, and assayed for luminescence in a luminometer which added 100  $\mu$ l of 1 mM luciferin.

30 Human HeLa cells (ATCC) were grown in RPMI 1640-10% fetal calf serum in an atmosphere of 5%  $CO_2$ . For infection of dividing cultures, cells were seeded at a density of  $1.6 \times 10^5$ /well in a 6-well tray the day before infection. G1/S arrested cultures were prepared seeding  $2 \times 10^5$  cells/well two days before infection, and adding 15  $\mu$ g/ml aphidicolin 24 hrs before infection. Aphidicolin was added daily to the medium throughout the infection and post-infection time, as described. G2-arrested cells were prepared by exposing the cells for 20 min to a  $^{60}Co$  source

calibrated at 200 rad/min one day before infection and seeding the cells at  $4 \times 10^5$ /well. Cell cycle arrest at the indicating stage of the cycle was confirmed by propidium iodide staining and flow cytometry.

5 Figure 4 shows the relative efficiency of transduction of CMV- $\beta$ -Gal into HeLa cells. The cells were growth arrested using either pharmacological means (aphidicolin) or x-irradiation. The infection rate for the HIV-based vector was more efficient when cells were arrested at the G1/S interphase by aphidicolin, however, the HIV-based vector was more efficient than MLV-based vector overall.

10 Rat 208F fibroblasts (a gift of B. Sefton, Salk Institute, La Jolla, CA) were grown in DMEM-10% calf serum in an atmosphere of 10% CO<sub>2</sub>. For infection of growing cultures, cells were seeded at a density of  $10^5$  cells/well in a 6-well tray the day before infection. Growth-arrested cultures were prepared by seeding  $2.5 \times 10^5$  cells/well in a 6-well tissue culture tray, and shifting the cultures to medium containing 5% calf serum and 2  $\mu$ M dexamethazone after they reached confluence, as described by  
15 Miller *et al.* (*Mol. Cell. Biol.*, 10:4239, 1990). Medium was changed every three-four days, and the cultures were kept for two to four weeks. When not otherwise indicated, G<sub>0</sub> cultures were used 3 weeks after reaching confluence. Growth arrest was documented by propidium iodide staining and flow cytometry. The fraction of cells in S-phase was 40 to 50% in growing cells, and from 10 down to 2% after reaching  
20 confluence, depending on the time past it. For rescue experiments, infected G<sub>0</sub> cultures were trypsinized at the indicated time after infection, and the cells replated at several dilutions. Expression of  $\beta$ -Gal was scored 48 hrs from replating.

Figure 5 shows the relative efficiency of transduction of CMV-luciferase into rat 208F fibroblasts with the HIV- and MLV-based vectors in cells growth arrested for 4, 7, 11  
25 or 15 days. The results show that the vectors both efficiently infect growing cells. The efficiency of transduction of the HIV-based vector versus the MLV-based vector was approximately 4-10 fold higher, depending on the length of time of the growth arrest.

Human peripheral blood monocytes were prepared from the buffy coats of healthy donors as previously described (von Schwedler *et al.*, *Proc. Natl. Acad. Sci. USA*,  
30 91:6992, 1994). Monocyte-derived macrophages were cultured in RPM1-10% human serum (Sigma) for two to four weeks prior to infection. Figure 6 shows the results of transduction of CMV-luciferase into human primary macrophages using the HIV-based and MLV-based vectors, either with or without an envelope. The MLV-based

-21-

vector was almost completely ineffective in infecting the cells and was similar in infectivity to the non-enveloped virus. In contrast, the HIV-based vector was highly efficient in transducing primary macrophages.

Further, the survival of the HIV-based vector was assessed in growth arrested cells that were passaged over a period of 2-8 days. Survival of the MLV-based vector was very low in cells at G0 and after replating at 2, 4, and 8 days. However, the HIV-based vector maintained at about 45-50% infectivity in the same cells. While not wanting to be bound by a particular theory, it is believed that HIV can remain extrachromosomal without being degraded, unlike MLV. Therefore, HIV is stable without integration.

#### Example 7

#### INFECTION *in vivo*

For *in vivo* experiments, HIV-based and MLV-based vectors carrying a CMV-LacZ marker gene were used. Both vector types were pseudotyped with VSV.G envelope proteins. Vectors were purified and concentrated to a titer of  $3 \times 10^8$  I.U./ml in a vehicle of sterile PBS supplemented with 2 µg/ml polybrene. All procedures were performed according to institution-approved protocols for animal work, and in a biosafety level 3 (BSL3) environment. Normal adult female Fischer rats were anesthetized with a mixture of ketamine (44 mg/kg), acepromazine (0.75 mg/kg) and xylozine (4 mg/kg) in 0.9% NaCl. 2 µl of viral suspension were slowly injected in the corpus striatum of both sides of the brain with a stereotactical guide using a 5 µl Hamilton syringe with a 26 gauge beveled needle. 7 days after injection the rats were deeply anesthetized and perfused with 4% formaldehyde/0.1% glutaraldehyde. The brains were removed and free-floating coronal freezing microtome sections (40 µm) were processed in parallel series of one section out of every five cut. One series was processed for β-Gal immunocytochemistry with a monoclonal antibody and stained with avidin/biotin peroxidase and diaminobenzidine/NiCl<sub>2</sub> as substrate. Another series was processed for triple immunofluorescence with β-Gal (rabbit), NeuN (monoclonal), and GFAP (guinea pig) antibodies using FITC-, Texas Red-, or Cy5-labeled secondary antibodies. NeuN is a nuclear marker reported to be present in terminally differentiated neurons (Mullen, *Devel.op.*, 116:201, 1992). Glial fibrillar acidic protein (GFAP) is a astrocyte differentiation marker. The sections were then analyzed by confocal scanning laser microscopy (bioRad MRT600). Immunofluorescent signals were collected, digitally color enhanced and superimposed. False color images were generated electronically using Adobe Photoshop (Adobe System Inc.).

-22-

The results revealed numerous  $\beta$ -gal positive neurons in regions injected with the HIV-based vector, but not in those inoculated with the MLV-based vector.

### SUMMARY

5 The results indicated the feasibility of gene transfer using a lentiviral vector. Using growing cells, the transduction efficiency obtained with a three-plasmid, transient packaging system was comparable to that obtained with similarly assembled, conventional (MLV-based) retroviral vectors. In sharp contrast with MLV-based vectors, the lentiviral vector was capable of efficient gene transfer in the absence of division of the target cell. Efficiency of transduction into non-dividing cells *in vitro* was dependent on the stage of cell cycle arrest. Infection of G1/S and G2 arrested cells was as efficient as that observed with growing cells. Infection of cells arrested in G<sub>0</sub> 10 *in vitro* was less efficient, the lesser the longer the cultures had been in G<sub>0</sub>. However, in long-term G<sub>0</sub> cultures the vector survived as a stable intermediate, and it could be rescued to complete its infectious cycle by stimulation of the cells to divide. Transduction of the marker gene into terminally differentiated cells, such as human 15 primary macrophages *in vitro* and rat cerebral neurons *in vivo*, was observed.

Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by 20 the following claims.



What is claimed is:

1. A recombinant retrovirus capable of infecting a non-dividing cell comprising:
  - a viral GAG;
  - a viral POL;
  - a viral ENV;
  - a heterologous nucleic acid sequence operably linked to a regulatory nucleic acid sequence; and
  - cis-acting nucleic acid sequences necessary for reverse transcription and integration.
2. The retrovirus of claim 1, wherein the virus is a lentivirus.
3. The retrovirus of claim 2, wherein the lentivirus is human immunodeficiency virus (HIV).
4. The retrovirus of claim 1, wherein the ENV is selected from the group consisting of Moloney leukemia virus (MLV) and Vesicular stomatitis virus (VSV) ENV.
5. The retrovirus of claim 1, wherein the ENV is amphotropic.
6. The retrovirus of claim 1, wherein the ENV is ecotropic.
7. The retrovirus of claim 1, further comprising a viral protein selected from the group consisting of VPR, VIF, NEF, VPX, TAT, REV, and VPU.
8. The retrovirus of claim 1, wherein the non-dividing cell is selected from the group consisting of a neuron, muscle, heart, lung, liver, skin, and bone marrow cell.
9. The retrovirus of claim 1, wherein the regulatory nucleic acid sequence is a promoter or an enhancer.
10. The retrovirus of claim 9, wherein the promoter is a viral promoter.

-24-

11. A method of producing a recombinant retrovirus capable of infecting a non-dividing cell, said method comprising:
  - transfecting a suitable packaging host cell with the following:
    - a vector providing a nucleic acid encoding a viral gag and a viral pol;
    - a vector providing a nucleic acid encoding a viral env;
    - a vector providing a nucleic acid sequence encoding a packaging signal flanked by cis-acting nucleic acid sequences for reverse transcription and integration, and a cloning site for introduction of a heterologous nucleic acid sequence, operably linked to a regulatory nucleic acid sequence; and
  - recovering the recombinant virus.
12. The method of claim 11, wherein the virus is a lentivirus.
13. The method of claim 12, wherein the lentivirus is human immunodeficiency virus (HIV).
14. The method of claim 11, wherein the transfecting further includes at least one vector providing a nucleic acid sequence selected from the group consisting of vpr, vif, nef, vpx, tat, rev, and vpu.
15. A recombinant retrovirus produced by the method of claim 11.
16. A method for introduction and expression of a heterologous nucleic acid sequence in a non-dividing cell comprising infecting the non-dividing cell with the recombinant virus of claim 1 and expressing the heterologous nucleic acid sequence in the non-dividing cell.
17. The method of claim 16, wherein the heterologous nucleic acid is an antisense nucleic acid.
18. The method of claim 16, wherein the heterologous nucleic acid is a biological modifier nucleic acid.

-25-

19. The method of claim 16, wherein the non-dividing cell is selected from the group consisting of a neuron, muscle, liver, skin, heart, lung, and bone marrow cell.
20. The method of claim 16, wherein the introduction is *ex vivo*.
21. The method of claim 16, wherein the introduction is *in vivo*.
22. The method of claim 16, wherein the virus is a lentivirus.

1 / 7




FIG. 1

2 / 7

TRANSFECTANT	INFECTIOUS TITER ON 208F FIBROBLASTS I.U./ml		YIELD
	CONDITIONED MEDIUM	PURIFIED STOCK	
pCMVDR8/ pHR'CLacZ/VSV.G	$6 \cdot 10^5$	$3.5 \cdot 10^8$	30%
PCMVGAGPOL(MLV) pSLX-CLacZ/VSV.G	$10^5$	$3 \cdot 10^8$	25%

FIG. 2

3 / 7

CO-TRANSFECTION IN 293T CELLS				
PACKAGING CONSTRUCT	PLASMID CARRYING CMV-LacZ	Env-ENCODING PLASMID	INFECTION OF RAT 208F CELLS I.U./ml	
HIV-BASED		pHR'	MLV (Ampho)	0 (<1/ml)
	pCMVDR8	pHR'	-	0
	pCMVDR8	pHR'	MLV (Ampho)	10 <sup>5</sup>
	pCMVDR8	pHR'	MLV (Eco)	10 <sup>4</sup>
	pCMVDR8	pHR'	VSV.G	5 10 <sup>5</sup>
	pCMVDR8	pLNL-SLX	MLV (Ampho)	0
	pCMVDR8	pGEM	MLV (Ampho)	0
MLV-BASED	pCL-ECO	pLNL-SLX		10 <sup>5</sup>
	pCL-ECO	pHR'		0
	pCMVGAGPOL	pLNL-SLX	MLV (Ampho)	10 <sup>5</sup>

FIG. 3

4 / 7




FIG. 4

SUBSTITUTE SHEET (RULE 26)

5/7




FIG. 5



6 / 7




FIG. 6

SUBSTITUTE SHEET (RULE 26)

7/7




FIG. 7

SUBSTITUTE SHEET (RULE 26)

## INTERNATIONAL SEARCH REPORT

national application No.  
PCT/US96/15406**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A61K 35/76, 48/00

US CL : 424/93.6; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/93.6; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
APS, MEDLINE, BIOSIS, CAPLUS, DISSABS, DERWENT, JAPIO, WIPO**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	LANDAU, N.R., et al. Pseudotyping with human T-cell leukemia virus type I broadens the human immunodeficiency virus host range. Journal of Virology. January 1991, Vol. 65, No. 1, pages 162-169, see entire document.	1-10 ----- 11-22

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

24 DECEMBER 1996

Date of mailing of the international search report

15 JAN 1997

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

PATRICK TWOMEY, PH.D.

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196